

# Background

To begin, recall that (somewhat informally) a **set** is simply a collection of objects (I encourage you to review the logic/proof notes for further details).

# Background

To begin, recall that (somewhat informally) a **set** is simply a collection of objects (I encourage you to review the logic/proof notes for further details). For example,  $\{1, 2, 3\}$  denotes the set whose **members** are the integers 1, 2, and 3.

# Background

To begin, recall that (somewhat informally) a **set** is simply a collection of objects (I encourage you to review the logic/proof notes for further details). For example,  $\{1, 2, 3\}$  denotes the set whose **members** are the integers 1, 2, and 3. When a pattern emerges, we often use an ellipsis to denote this.

# Background

To begin, recall that (somewhat informally) a **set** is simply a collection of objects (I encourage you to review the logic/proof notes for further details). For example,  $\{1, 2, 3\}$  denotes the set whose **members** are the integers 1, 2, and 3. When a pattern emerges, we often use an ellipsis to denote this. For instance,  $\{1, 2, 3, 4, \dots\}$  denotes the set of *all* positive integers.

# Background

To begin, recall that (somewhat informally) a **set** is simply a collection of objects (I encourage you to review the logic/proof notes for further details). For example,  $\{1, 2, 3\}$  denotes the set whose **members** are the integers 1, 2, and 3. When a pattern emerges, we often use an ellipsis to denote this. For instance,  $\{1, 2, 3, 4, \dots\}$  denotes the set of *all* positive integers. Sometimes sets are sufficiently complicated as to necessitate other methods of describing them.

# Background

To begin, recall that (somewhat informally) a **set** is simply a collection of objects (I encourage you to review the logic/proof notes for further details). For example,  $\{1, 2, 3\}$  denotes the set whose **members** are the integers 1, 2, and 3. When a pattern emerges, we often use an ellipsis to denote this. For instance,  $\{1, 2, 3, 4, \dots\}$  denotes the set of *all* positive integers. Sometimes sets are sufficiently complicated as to necessitate other methods of describing them. For example,  $\{n: n \text{ is a prime number}\}$  denotes the set of all objects  $n$  with the property that  $n$  is a prime number.

# Background

To begin, recall that (somewhat informally) a **set** is simply a collection of objects (I encourage you to review the logic/proof notes for further details). For example,  $\{1, 2, 3\}$  denotes the set whose **members** are the integers 1, 2, and 3. When a pattern emerges, we often use an ellipsis to denote this. For instance,  $\{1, 2, 3, 4, \dots\}$  denotes the set of *all* positive integers. Sometimes sets are sufficiently complicated as to necessitate other methods of describing them. For example,  $\{n: n \text{ is a prime number}\}$  denotes the set of all objects  $n$  with the property that  $n$  is a prime number. Since the following sets of numbers will come up often in this course, let's review the notation.

# Background

To begin, recall that (somewhat informally) a **set** is simply a collection of objects (I encourage you to review the logic/proof notes for further details). For example,  $\{1, 2, 3\}$  denotes the set whose **members** are the integers 1, 2, and 3. When a pattern emerges, we often use an ellipsis to denote this. For instance,  $\{1, 2, 3, 4, \dots\}$  denotes the set of *all* positive integers. Sometimes sets are sufficiently complicated as to necessitate other methods of describing them. For example,  $\{n: n \text{ is a prime number}\}$  denotes the set of all objects  $n$  with the property that  $n$  is a prime number. Since the following sets of numbers will come up often in this course, let's review the notation.

# Background

## Definition

Recall the following:

# Background

## Definition

Recall the following:

- 1  $\mathbb{N}$ , the set of **natural numbers**, is given by  $\{0, 1, 2, 3, \dots\}$ ,

# Background

## Definition

Recall the following:

- 1  $\mathbb{N}$ , the set of **natural numbers**, is given by  $\{0, 1, 2, 3, \dots\}$ ,
- 2  $\mathbb{Z}$ , the set of **integers**, is given by  $\{\dots, -3, -2, -1, 0, 1, 2, 3, \dots\}$ ,

# Background

## Definition

Recall the following:

- 1  $\mathbb{N}$ , the set of **natural numbers**, is given by  $\{0, 1, 2, 3, \dots\}$ ,
- 2  $\mathbb{Z}$ , the set of **integers**, is given by  $\{\dots, -3, -2, -1, 0, 1, 2, 3, \dots\}$ ,
- 3  $\mathbb{Q}$ , the set of **rational numbers**, is given by  $\{x: x = \frac{p}{q} \text{ for some integers } p \text{ and } q \text{ with } q \neq 0\}$ .

# Background

## Definition

Recall the following:

- 1  $\mathbb{N}$ , the set of **natural numbers**, is given by  $\{0, 1, 2, 3, \dots\}$ ,
- 2  $\mathbb{Z}$ , the set of **integers**, is given by  $\{\dots, -3, -2, -1, 0, 1, 2, 3, \dots\}$ ,
- 3  $\mathbb{Q}$ , the set of **rational numbers**, is given by  $\{x: x = \frac{p}{q} \text{ for some integers } p \text{ and } q \text{ with } q \neq 0\}$ .
- 4  $\mathbb{R}$ , the set of **real numbers**, is (informally) the set of all numbers you've encountered which do not have an "i" in them (it is possible to give a formal definition of this set of numbers, but it involves a lot of axiomatic set theory which is beyond the scope of this course).

# Background

## Definition

Recall the following:

- 1  $\mathbb{N}$ , the set of **natural numbers**, is given by  $\{0, 1, 2, 3, \dots\}$ ,
- 2  $\mathbb{Z}$ , the set of **integers**, is given by  $\{\dots, -3, -2, -1, 0, 1, 2, 3, \dots\}$ ,
- 3  $\mathbb{Q}$ , the set of **rational numbers**, is given by  $\{x: x = \frac{p}{q} \text{ for some integers } p \text{ and } q \text{ with } q \neq 0\}$ .
- 4  $\mathbb{R}$ , the set of **real numbers**, is (informally) the set of all numbers you've encountered which do not have an "i" in them (it is possible to give a formal definition of this set of numbers, but it involves a lot of axiomatic set theory which is beyond the scope of this course).
- 5  $\mathbb{C}$ , the set of **complex numbers**, is given by  $\{x: x = a + bi \text{ for some real numbers } a \text{ and } b\}$ , where  $i^2 = -1$ .

# Background

## Definition

Recall the following:

- 1  $\mathbb{N}$ , the set of **natural numbers**, is given by  $\{0, 1, 2, 3, \dots\}$ ,
- 2  $\mathbb{Z}$ , the set of **integers**, is given by  $\{\dots, -3, -2, -1, 0, 1, 2, 3, \dots\}$ ,
- 3  $\mathbb{Q}$ , the set of **rational numbers**, is given by  $\{x: x = \frac{p}{q} \text{ for some integers } p \text{ and } q \text{ with } q \neq 0\}$ .
- 4  $\mathbb{R}$ , the set of **real numbers**, is (informally) the set of all numbers you've encountered which do not have an "i" in them (it is possible to give a formal definition of this set of numbers, but it involves a lot of axiomatic set theory which is beyond the scope of this course).
- 5  $\mathbb{C}$ , the set of **complex numbers**, is given by  $\{x: x = a + bi \text{ for some real numbers } a \text{ and } b\}$ , where  $i^2 = -1$ .

# Operations

It is important to note that in mathematics, we often will consider sets of object which are NOT numbers.

# Operations

It is important to note that in mathematics, we often will consider sets of object which are NOT numbers. For example, we can talk about a set of geometric figures, sets of functions, sets of matrices, and so on.

# Operations

It is important to note that in mathematics, we often will consider sets of object which are NOT numbers. For example, we can talk about a set of geometric figures, sets of functions, sets of matrices, and so on. So please do NOT assume members of sets are always numbers.

# Operations

It is important to note that in mathematics, we often will consider sets of object which are NOT numbers. For example, we can talk about a set of geometric figures, sets of functions, sets of matrices, and so on. So please do NOT assume members of sets are always numbers.

Now that we have tersely reviewed sets, we will be interested in defining so-called *operations* on sets, defined as follows:

# Operations

It is important to note that in mathematics, we often will consider sets of object which are NOT numbers. For example, we can talk about a set of geometric figures, sets of functions, sets of matrices, and so on. So please do NOT assume members of sets are always numbers.

Now that we have tersely reviewed sets, we will be interested in defining so-called *operations* on sets, defined as follows:

## Definition

Let  $S$  be a set (remember,  $S$  need not contain numbers; view  $S$  as a collection of objects but that you do not know what the objects are).

# Operations

It is important to note that in mathematics, we often will consider sets of object which are NOT numbers. For example, we can talk about a set of geometric figures, sets of functions, sets of matrices, and so on. So please do NOT assume members of sets are always numbers.

Now that we have tersely reviewed sets, we will be interested in defining so-called *operations* on sets, defined as follows:

## Definition

Let  $S$  be a set (remember,  $S$  need not contain numbers; view  $S$  as a collection of objects but that you do not know what the objects are). An **operation** on  $S$  is a rule  $*$  for combining

# Operations

It is important to note that in mathematics, we often will consider sets of object which are NOT numbers. For example, we can talk about a set of geometric figures, sets of functions, sets of matrices, and so on. So please do NOT assume members of sets are always numbers.

Now that we have tersely reviewed sets, we will be interested in defining so-called *operations* on sets, defined as follows:

## Definition

Let  $S$  be a set (remember,  $S$  need not contain numbers; view  $S$  as a collection of objects but that you do not know what the objects are). An **operation** on  $S$  is a rule  $*$  for combining

- 1 any members  $a$  and  $b$  of  $S$  (possibly  $a = b$ ) to produce

# Operations

It is important to note that in mathematics, we often will consider sets of object which are NOT numbers. For example, we can talk about a set of geometric figures, sets of functions, sets of matrices, and so on. So please do NOT assume members of sets are always numbers.

Now that we have tersely reviewed sets, we will be interested in defining so-called *operations* on sets, defined as follows:

## Definition

Let  $S$  be a set (remember,  $S$  need not contain numbers; view  $S$  as a collection of objects but that you do not know what the objects are). An **operation** on  $S$  is a rule  $*$  for combining

- 1 any members  $a$  and  $b$  of  $S$  (possibly  $a = b$ ) to produce
- 2 a *unique* (i.e. exactly one) member of  $S$ .

# Operations

It is important to note that in mathematics, we often will consider sets of object which are NOT numbers. For example, we can talk about a set of geometric figures, sets of functions, sets of matrices, and so on. So please do NOT assume members of sets are always numbers.

Now that we have tersely reviewed sets, we will be interested in defining so-called *operations* on sets, defined as follows:

## Definition

Let  $S$  be a set (remember,  $S$  need not contain numbers; view  $S$  as a collection of objects but that you do not know what the objects are). An **operation** on  $S$  is a rule  $*$  for combining

- 1 any members  $a$  and  $b$  of  $S$  (possibly  $a = b$ ) to produce
- 2 a *unique* (i.e. exactly one) member of  $S$ .

# Operations

## Example

Suppose that  $S = \{0, 1\}$  and define  $*$  on  $S$  by declaring that  $0 * 0 = 0$ ,  $0 * 1 = 0$ ,  $1 * 0 = 0$ , and  $1 * 1 = 1$ .

# Operations

## Example

Suppose that  $S = \{0, 1\}$  and define  $*$  on  $S$  by declaring that  $0 * 0 = 0$ ,  $0 * 1 = 0$ ,  $1 * 0 = 0$ , and  $1 * 1 = 1$ . Observe that  $*$  combines *any* two members of  $S$  and upon combining, we get *exactly one* member of  $S$

# Operations

## Example

Suppose that  $S = \{0, 1\}$  and define  $*$  on  $S$  by declaring that  $0 * 0 = 0$ ,  $0 * 1 = 0$ ,  $1 * 0 = 0$ , and  $1 * 1 = 1$ . Observe that  $*$  combines *any* two members of  $S$  and upon combining, we get *exactly one* member of  $S$  (note that in this case,  $*$  is simply the usual multiplication on  $S$ ).

# Operations

## Example

Suppose that  $S = \{0, 1\}$  and define  $*$  on  $S$  by declaring that  $0 * 0 = 0$ ,  $0 * 1 = 0$ ,  $1 * 0 = 0$ , and  $1 * 1 = 1$ . Observe that  $*$  combines *any* two members of  $S$  and upon combining, we get *exactly one* member of  $S$  (note that in this case,  $*$  is simply the usual multiplication on  $S$ ).

## Example

Let  $S$  be the set  $\mathbb{R}$  of real numbers.

# Operations

## Example

Suppose that  $S = \{0, 1\}$  and define  $*$  on  $S$  by declaring that  $0 * 0 = 0$ ,  $0 * 1 = 0$ ,  $1 * 0 = 0$ , and  $1 * 1 = 1$ . Observe that  $*$  combines *any* two members of  $S$  and upon combining, we get *exactly one* member of  $S$  (note that in this case,  $*$  is simply the usual multiplication on  $S$ ).

## Example

Let  $S$  be the set  $\mathbb{R}$  of real numbers. Now define  $*$  on  $S$  by  $a * b = \min(a, b)$ , where  $\min(a, b)$  denotes the smaller of the numbers  $a$  and  $b$ .

# Operations

## Example

Suppose that  $S = \{0, 1\}$  and define  $*$  on  $S$  by declaring that  $0 * 0 = 0$ ,  $0 * 1 = 0$ ,  $1 * 0 = 0$ , and  $1 * 1 = 1$ . Observe that  $*$  combines *any* two members of  $S$  and upon combining, we get *exactly one* member of  $S$  (note that in this case,  $*$  is simply the usual multiplication on  $S$ ).

## Example

Let  $S$  be the set  $\mathbb{R}$  of real numbers. Now define  $*$  on  $S$  by  $a * b = \min(a, b)$ , where  $\min(a, b)$  denotes the smaller of the numbers  $a$  and  $b$ . In case  $a = b$ , we declare  $\min(a, b) = a$ .

# Operations

## Example

Suppose that  $S = \{0, 1\}$  and define  $*$  on  $S$  by declaring that  $0 * 0 = 0$ ,  $0 * 1 = 0$ ,  $1 * 0 = 0$ , and  $1 * 1 = 1$ . Observe that  $*$  combines *any* two members of  $S$  and upon combining, we get *exactly one* member of  $S$  (note that in this case,  $*$  is simply the usual multiplication on  $S$ ).

## Example

Let  $S$  be the set  $\mathbb{R}$  of real numbers. Now define  $*$  on  $S$  by  $a * b = \min(a, b)$ , where  $\min(a, b)$  denotes the smaller of the numbers  $a$  and  $b$ . In case  $a = b$ , we declare  $\min(a, b) = a$ . Then  $*$  is defined for *any* real numbers  $a$  and  $b$  and results in a *unique* real number

# Operations

## Example

Suppose that  $S = \{0, 1\}$  and define  $*$  on  $S$  by declaring that  $0 * 0 = 0$ ,  $0 * 1 = 0$ ,  $1 * 0 = 0$ , and  $1 * 1 = 1$ . Observe that  $*$  combines *any* two members of  $S$  and upon combining, we get *exactly one* member of  $S$  (note that in this case,  $*$  is simply the usual multiplication on  $S$ ).

## Example

Let  $S$  be the set  $\mathbb{R}$  of real numbers. Now define  $*$  on  $S$  by  $a * b = \min(a, b)$ , where  $\min(a, b)$  denotes the smaller of the numbers  $a$  and  $b$ . In case  $a = b$ , we declare  $\min(a, b) = a$ . Then  $*$  is defined for *any* real numbers  $a$  and  $b$  and results in a *unique* real number (note that, for example  $\min(1, 2) = 1$ ;

# Operations

## Example

Suppose that  $S = \{0, 1\}$  and define  $*$  on  $S$  by declaring that  $0 * 0 = 0$ ,  $0 * 1 = 0$ ,  $1 * 0 = 0$ , and  $1 * 1 = 1$ . Observe that  $*$  combines *any* two members of  $S$  and upon combining, we get *exactly one* member of  $S$  (note that in this case,  $*$  is simply the usual multiplication on  $S$ ).

## Example

Let  $S$  be the set  $\mathbb{R}$  of real numbers. Now define  $*$  on  $S$  by  $a * b = \min(a, b)$ , where  $\min(a, b)$  denotes the smaller of the numbers  $a$  and  $b$ . In case  $a = b$ , we declare  $\min(a, b) = a$ . Then  $*$  is defined for *any* real numbers  $a$  and  $b$  and results in a *unique* real number (note that, for example  $\min(1, 2) = 1$ ; the min is not equal to both 1 and 4, for example; this would violate uniqueness).

# Operations

## Example

Suppose that  $S = \{0, 1\}$  and define  $*$  on  $S$  by declaring that  $0 * 0 = 0$ ,  $0 * 1 = 0$ ,  $1 * 0 = 0$ , and  $1 * 1 = 1$ . Observe that  $*$  combines *any* two members of  $S$  and upon combining, we get *exactly one* member of  $S$  (note that in this case,  $*$  is simply the usual multiplication on  $S$ ).

## Example

Let  $S$  be the set  $\mathbb{R}$  of real numbers. Now define  $*$  on  $S$  by  $a * b = \min(a, b)$ , where  $\min(a, b)$  denotes the smaller of the numbers  $a$  and  $b$ . In case  $a = b$ , we declare  $\min(a, b) = a$ . Then  $*$  is defined for *any* real numbers  $a$  and  $b$  and results in a *unique* real number (note that, for example  $\min(1, 2) = 1$ ; the min is not equal to both 1 and 4, for example; this would violate uniqueness).

# Operations

## Example

Let  $S = \mathbb{R}$ , and define  $a * b = \sqrt{ab}$ .

# Operations

## Example

Let  $S = \mathbb{R}$ , and define  $a * b = \sqrt{ab}$ . Is  $*$  an operation on  $S$ ?

# Operations

## Example

Let  $S = \mathbb{R}$ , and define  $a * b = \sqrt{ab}$ . Is  $*$  an operation on  $S$ ?

*Solution.* No,  $*$  is not an operation on  $S$ .

# Operations

## Example

Let  $S = \mathbb{R}$ , and define  $a * b = \sqrt{ab}$ . Is  $*$  an operation on  $S$ ?

*Solution.* No,  $*$  is not an operation on  $S$ . One way to see this is that  $1 * -1 = \sqrt{1 \cdot -1} = \sqrt{-1} = i \notin \mathbb{R}$ .

# Operations

## Example

Let  $S = \mathbb{R}$ , and define  $a * b = \sqrt{ab}$ . Is  $*$  an operation on  $S$ ?

*Solution.* No,  $*$  is not an operation on  $S$ . One way to see this is that  $1 * -1 = \sqrt{1 \cdot -1} = \sqrt{-1} = i \notin \mathbb{R}$ . Remember that to be an operation, when we combine *any* two elements of  $S$  we must get an element which is still in  $S$ .  
 $\square$

# Operations

## Example

Let  $S = \mathbb{R}$ , and define  $a * b = \sqrt{ab}$ . Is  $*$  an operation on  $S$ ?

*Solution.* No,  $*$  is not an operation on  $S$ . One way to see this is that  $1 * -1 = \sqrt{1 \cdot -1} = \sqrt{-1} = i \notin \mathbb{R}$ . Remember that to be an operation, when we combine *any* two elements of  $S$  we must get an element which is still in  $S$ .  
□

## Example

To get a bit more abstract, imagine that  $S$  contains all paints (of every color on the spectrum, let's say).

# Operations

## Example

Let  $S = \mathbb{R}$ , and define  $a * b = \sqrt{ab}$ . Is  $*$  an operation on  $S$ ?

*Solution.* No,  $*$  is not an operation on  $S$ . One way to see this is that  $1 * -1 = \sqrt{1 \cdot -1} = \sqrt{-1} = i \notin \mathbb{R}$ . Remember that to be an operation, when we combine *any* two elements of  $S$  we must get an element which is still in  $S$ .  
□

## Example

To get a bit more abstract, imagine that  $S$  contains all paints (of every color on the spectrum, let's say). We can define  $*$  on  $S$  by  $c_1 * c_2$  to mean mix (say equal amounts of) paint of color  $c_1$  and paint of color  $c_2$ .

# Operations

## Example

Let  $S = \mathbb{R}$ , and define  $a * b = \sqrt{ab}$ . Is  $*$  an operation on  $S$ ?

*Solution.* No,  $*$  is not an operation on  $S$ . One way to see this is that  $1 * -1 = \sqrt{1 \cdot -1} = \sqrt{-1} = i \notin \mathbb{R}$ . Remember that to be an operation, when we combine *any* two elements of  $S$  we must get an element which is still in  $S$ .  
□

## Example

To get a bit more abstract, imagine that  $S$  contains all paints (of every color on the spectrum, let's say). We can define  $*$  on  $S$  by  $c_1 * c_2$  to mean mix (say equal amounts of) paint of color  $c_1$  and paint of color  $c_2$ . Then you will get paint of some color (for example, mixing yellow and blue gives green).

# Operations

## Example

Let  $S = \mathbb{R}$ , and define  $a * b = \sqrt{ab}$ . Is  $*$  an operation on  $S$ ?

*Solution.* No,  $*$  is not an operation on  $S$ . One way to see this is that  $1 * -1 = \sqrt{1 \cdot -1} = \sqrt{-1} = i \notin \mathbb{R}$ . Remember that to be an operation, when we combine *any* two elements of  $S$  we must get an element which is still in  $S$ .  
□

## Example

To get a bit more abstract, imagine that  $S$  contains all paints (of every color on the spectrum, let's say). We can define  $*$  on  $S$  by  $c_1 * c_2$  to mean mix (say equal amounts of) paint of color  $c_1$  and paint of color  $c_2$ . Then you will get paint of some color (for example, mixing yellow and blue gives green).

# Operations

## Example

Let  $S = \mathbb{N}$  and define  $*$  on  $S$  by  $a * b = 0$  (that is,  $a * b = 0$  for EVERY natural number  $a$  and  $b$ ).

# Operations

## Example

Let  $S = \mathbb{N}$  and define  $*$  on  $S$  by  $a * b = 0$  (that is,  $a * b = 0$  for EVERY natural number  $a$  and  $b$ ). Is  $*$  an operation?

# Operations

## Example

Let  $S = \mathbb{N}$  and define  $*$  on  $S$  by  $a * b = 0$  (that is,  $a * b = 0$  for EVERY natural number  $a$  and  $b$ ). Is  $*$  an operation?

*Solution.* Yes,  $*$  is an operation.

# Operations

## Example

Let  $S = \mathbb{N}$  and define  $*$  on  $S$  by  $a * b = 0$  (that is,  $a * b = 0$  for EVERY natural number  $a$  and  $b$ ). Is  $*$  an operation?

*Solution.* Yes,  $*$  is an operation. First, for any natural numbers  $a$  and  $b$ , I have defined  $a * b$  to be 0,

# Operations

## Example

Let  $S = \mathbb{N}$  and define  $*$  on  $S$  by  $a * b = 0$  (that is,  $a * b = 0$  for EVERY natural number  $a$  and  $b$ ). Is  $*$  an operation?

*Solution.* Yes,  $*$  is an operation. First, for any natural numbers  $a$  and  $b$ , I have defined  $a * b$  to be 0, so the rule is defined for every natural numbers  $a$  and  $b$  and results again in a natural number.

# Operations

## Example

Let  $S = \mathbb{N}$  and define  $*$  on  $S$  by  $a * b = 0$  (that is,  $a * b = 0$  for EVERY natural number  $a$  and  $b$ ). Is  $*$  an operation?

*Solution.* Yes,  $*$  is an operation. First, for any natural numbers  $a$  and  $b$ , I have defined  $a * b$  to be 0, so the rule is defined for every natural numbers  $a$  and  $b$  and results again in a natural number. Now (this is important), for any natural numbers  $a$  and  $b$ ,  $a * b$  is uniquely determined:

# Operations

## Example

Let  $S = \mathbb{N}$  and define  $*$  on  $S$  by  $a * b = 0$  (that is,  $a * b = 0$  for EVERY natural number  $a$  and  $b$ ). Is  $*$  an operation?

*Solution.* Yes,  $*$  is an operation. First, for any natural numbers  $a$  and  $b$ , I have defined  $a * b$  to be 0, so the rule is defined for every natural numbers  $a$  and  $b$  and results again in a natural number. Now (this is important), for any natural numbers  $a$  and  $b$ ,  $a * b$  is uniquely determined: it is 0 (not, for example 0 and 1 and 5; THAT would violate uniqueness). □

# Operations

## Example

Let  $S = \mathbb{N}$  and define  $*$  on  $S$  by  $a * b = 0$  (that is,  $a * b = 0$  for EVERY natural number  $a$  and  $b$ ). Is  $*$  an operation?

*Solution.* Yes,  $*$  is an operation. First, for any natural numbers  $a$  and  $b$ , I have defined  $a * b$  to be 0, so the rule is defined for every natural numbers  $a$  and  $b$  and results again in a natural number. Now (this is important), for any natural numbers  $a$  and  $b$ ,  $a * b$  is uniquely determined: it is 0 (not, for example 0 and 1 and 5; THAT would violate uniqueness). □

## Example

Let  $S$  be the set of positive real numbers. Is  $a * b = \pm\sqrt{ab}$  an operation on  $S$ ?

# Operations

## Example

Let  $S = \mathbb{N}$  and define  $*$  on  $S$  by  $a * b = 0$  (that is,  $a * b = 0$  for EVERY natural number  $a$  and  $b$ ). Is  $*$  an operation?

*Solution.* Yes,  $*$  is an operation. First, for any natural numbers  $a$  and  $b$ , I have defined  $a * b$  to be 0, so the rule is defined for every natural numbers  $a$  and  $b$  and results again in a natural number. Now (this is important), for any natural numbers  $a$  and  $b$ ,  $a * b$  is uniquely determined: it is 0 (not, for example 0 and 1 and 5; THAT would violate uniqueness). □

## Example

Let  $S$  be the set of positive real numbers. Is  $a * b = \pm\sqrt{ab}$  an operation on  $S$ ?

*Solution.* No.

# Operations

## Example

Let  $S = \mathbb{N}$  and define  $*$  on  $S$  by  $a * b = 0$  (that is,  $a * b = 0$  for EVERY natural number  $a$  and  $b$ ). Is  $*$  an operation?

*Solution.* Yes,  $*$  is an operation. First, for any natural numbers  $a$  and  $b$ , I have defined  $a * b$  to be 0, so the rule is defined for every natural numbers  $a$  and  $b$  and results again in a natural number. Now (this is important), for any natural numbers  $a$  and  $b$ ,  $a * b$  is uniquely determined: it is 0 (not, for example 0 and 1 and 5; THAT would violate uniqueness).  $\square$

## Example

Let  $S$  be the set of positive real numbers. Is  $a * b = \pm\sqrt{ab}$  an operation on  $S$ ?

*Solution.* No. This definition violates both parts of the definition.

# Operations

## Example

Let  $S = \mathbb{N}$  and define  $*$  on  $S$  by  $a * b = 0$  (that is,  $a * b = 0$  for EVERY natural number  $a$  and  $b$ ). Is  $*$  an operation?

*Solution.* Yes,  $*$  is an operation. First, for any natural numbers  $a$  and  $b$ , I have defined  $a * b$  to be 0, so the rule is defined for every natural numbers  $a$  and  $b$  and results again in a natural number. Now (this is important), for any natural numbers  $a$  and  $b$ ,  $a * b$  is uniquely determined: it is 0 (not, for example 0 and 1 and 5; THAT would violate uniqueness).  $\square$

## Example

Let  $S$  be the set of positive real numbers. Is  $a * b = \pm\sqrt{ab}$  an operation on  $S$ ?

*Solution.* No. This definition violates both parts of the definition. First, note that  $1 * 1 = \pm\sqrt{1 \cdot 1} = \pm\sqrt{1} = \pm 1$ .

# Operations

## Example

Let  $S = \mathbb{N}$  and define  $*$  on  $S$  by  $a * b = 0$  (that is,  $a * b = 0$  for EVERY natural number  $a$  and  $b$ ). Is  $*$  an operation?

*Solution.* Yes,  $*$  is an operation. First, for any natural numbers  $a$  and  $b$ , I have defined  $a * b$  to be 0, so the rule is defined for every natural numbers  $a$  and  $b$  and results again in a natural number. Now (this is important), for any natural numbers  $a$  and  $b$ ,  $a * b$  is uniquely determined: it is 0 (not, for example 0 and 1 and 5; THAT would violate uniqueness). □

## Example

Let  $S$  be the set of positive real numbers. Is  $a * b = \pm\sqrt{ab}$  an operation on  $S$ ?

*Solution.* No. This definition violates both parts of the definition. First, note that  $1 * 1 = \pm\sqrt{1 \cdot 1} = \pm\sqrt{1} = \pm 1$ . Thus  $1 * 1$  is NOT uniquely determined.

# Operations

## Example

Let  $S = \mathbb{N}$  and define  $*$  on  $S$  by  $a * b = 0$  (that is,  $a * b = 0$  for EVERY natural number  $a$  and  $b$ ). Is  $*$  an operation?

*Solution.* Yes,  $*$  is an operation. First, for any natural numbers  $a$  and  $b$ , I have defined  $a * b$  to be 0, so the rule is defined for every natural numbers  $a$  and  $b$  and results again in a natural number. Now (this is important), for any natural numbers  $a$  and  $b$ ,  $a * b$  is uniquely determined: it is 0 (not, for example 0 and 1 and 5; THAT would violate uniqueness). □

## Example

Let  $S$  be the set of positive real numbers. Is  $a * b = \pm\sqrt{ab}$  an operation on  $S$ ?

*Solution.* No. This definition violates both parts of the definition. First, note that  $1 * 1 = \pm\sqrt{1 \cdot 1} = \pm\sqrt{1} = \pm 1$ . Thus  $1 * 1$  is NOT uniquely determined. Moreover,  $-1$  is not in the set  $S$ . □

# Operations

## Example

Let  $S = \mathbb{N}$  and define  $*$  on  $S$  by  $a * b = 0$  (that is,  $a * b = 0$  for EVERY natural number  $a$  and  $b$ ). Is  $*$  an operation?

*Solution.* Yes,  $*$  is an operation. First, for any natural numbers  $a$  and  $b$ , I have defined  $a * b$  to be 0, so the rule is defined for every natural numbers  $a$  and  $b$  and results again in a natural number. Now (this is important), for any natural numbers  $a$  and  $b$ ,  $a * b$  is uniquely determined: it is 0 (not, for example 0 and 1 and 5; THAT would violate uniqueness). □

## Example

Let  $S$  be the set of positive real numbers. Is  $a * b = \pm\sqrt{ab}$  an operation on  $S$ ?

*Solution.* No. This definition violates both parts of the definition. First, note that  $1 * 1 = \pm\sqrt{1 \cdot 1} = \pm\sqrt{1} = \pm 1$ . Thus  $1 * 1$  is NOT uniquely determined. Moreover,  $-1$  is not in the set  $S$ . □

# Scalar Products

Next, we define the abstract concept of “scalar product.”

# Scalar Products

Next, we define the abstract concept of “scalar product.” For now, we will define this notion only relative to the set  $\mathbb{R}$  of real numbers.

# Scalar Products

Next, we define the abstract concept of “scalar product.” For now, we will define this notion only relative to the set  $\mathbb{R}$  of real numbers. This will get more abstract shortly.

## Definition

Let  $S$  be a set.

# Scalar Products

Next, we define the abstract concept of “scalar product.” For now, we will define this notion only relative to the set  $\mathbb{R}$  of real numbers. This will get more abstract shortly.

## Definition

Let  $S$  be a set. A **scalar product** on  $S$  (relative to  $\mathbb{R}$ ) is a rule  $\cdot$  for combining

# Scalar Products

Next, we define the abstract concept of “scalar product.” For now, we will define this notion only relative to the set  $\mathbb{R}$  of real numbers. This will get more abstract shortly.

## Definition

Let  $S$  be a set. A **scalar product** on  $S$  (relative to  $\mathbb{R}$ ) is a rule  $\cdot$  for combining

- 1 any real number  $r$  and any member  $s \in S$  to produce

# Scalar Products

Next, we define the abstract concept of “scalar product.” For now, we will define this notion only relative to the set  $\mathbb{R}$  of real numbers. This will get more abstract shortly.

## Definition

Let  $S$  be a set. A **scalar product** on  $S$  (relative to  $\mathbb{R}$ ) is a rule  $\cdot$  for combining

- 1 any real number  $r$  and any member  $s \in S$  to produce
- 2 a *unique* member of  $S$ .

# Scalar Products

Next, we define the abstract concept of “scalar product.” For now, we will define this notion only relative to the set  $\mathbb{R}$  of real numbers. This will get more abstract shortly.

## Definition

Let  $S$  be a set. A **scalar product** on  $S$  (relative to  $\mathbb{R}$ ) is a rule  $\cdot$  for combining

- 1 any real number  $r$  and any member  $s \in S$  to produce
- 2 a *unique* member of  $S$ .

Note the similarity to the definition of “operation”.

# Scalar Products

Next, we define the abstract concept of “scalar product.” For now, we will define this notion only relative to the set  $\mathbb{R}$  of real numbers. This will get more abstract shortly.

## Definition

Let  $S$  be a set. A **scalar product** on  $S$  (relative to  $\mathbb{R}$ ) is a rule  $\cdot$  for combining

- 1 any real number  $r$  and any member  $s \in S$  to produce
- 2 a *unique* member of  $S$ .

Note the similarity to the definition of “operation”. The main difference is that we are combining members of (in general) two different sets to produce a unique member of one of the sets.

# Scalar Products

Next, we define the abstract concept of “scalar product.” For now, we will define this notion only relative to the set  $\mathbb{R}$  of real numbers. This will get more abstract shortly.

## Definition

Let  $S$  be a set. A **scalar product** on  $S$  (relative to  $\mathbb{R}$ ) is a rule  $\cdot$  for combining

- 1 any real number  $r$  and any member  $s \in S$  to produce
- 2 a *unique* member of  $S$ .

Note the similarity to the definition of “operation”. The main difference is that we are combining members of (in general) two different sets to produce a unique member of one of the sets. As before, let’s look at some examples.

# Scalar Products

Next, we define the abstract concept of “scalar product.” For now, we will define this notion only relative to the set  $\mathbb{R}$  of real numbers. This will get more abstract shortly.

## Definition

Let  $S$  be a set. A **scalar product** on  $S$  (relative to  $\mathbb{R}$ ) is a rule  $\cdot$  for combining

- 1 any real number  $r$  and any member  $s \in S$  to produce
- 2 a *unique* member of  $S$ .

Note the similarity to the definition of “operation”. The main difference is that we are combining members of (in general) two different sets to produce a unique member of one of the sets. As before, let’s look at some examples.



# Scalar Products

## Example

Let  $S = \mathbb{R}^2 = \{(a, b) : a, b \in \mathbb{R}\}$  (that is,  $S$  is the set “ $\mathbb{R}^2$ ” of ordered pairs of real numbers; you should have worked with this set in calculus and a first course in linear algebra). Now define  $\cdot$  by  $r \cdot (a, b) = (ra, rb)$ , where  $ra$  and  $rb$  are just the usual product of the real numbers  $r$  and  $a$  and  $r$  and  $b$ , respectively.

# Scalar Products

## Example

Let  $S = \mathbb{R}^2 = \{(a, b) : a, b \in \mathbb{R}\}$  (that is,  $S$  is the set “ $\mathbb{R}^2$ ” of ordered pairs of real numbers; you should have worked with this set in calculus and a first course in linear algebra). Now define  $\cdot$  by  $r \cdot (a, b) = (ra, rb)$ , where  $ra$  and  $rb$  are just the usual product of the real numbers  $r$  and  $a$  and  $r$  and  $b$ , respectively. Then  $\cdot$  combines any real number with any member of  $\mathbb{R}^2$  to produce a unique member of  $\mathbb{R}^2$ .

# Scalar Products

## Example

Let  $S = \mathbb{R}^2 = \{(a, b) : a, b \in \mathbb{R}\}$  (that is,  $S$  is the set “ $\mathbb{R}^2$ ” of ordered pairs of real numbers; you should have worked with this set in calculus and a first course in linear algebra). Now define  $\cdot$  by  $r \cdot (a, b) = (ra, rb)$ , where  $ra$  and  $rb$  are just the usual product of the real numbers  $r$  and  $a$  and  $r$  and  $b$ , respectively. Then  $\cdot$  combines any real number with any member of  $\mathbb{R}^2$  to produce a unique member of  $\mathbb{R}^2$ . This is the so-called standard scalar product on the vector space  $\mathbb{R}^2$ .

# Scalar Products

## Example

Let  $S = \mathbb{R}^2 = \{(a, b) : a, b \in \mathbb{R}\}$  (that is,  $S$  is the set “ $\mathbb{R}^2$ ” of ordered pairs of real numbers; you should have worked with this set in calculus and a first course in linear algebra). Now define  $\cdot$  by  $r \cdot (a, b) = (ra, rb)$ , where  $ra$  and  $rb$  are just the usual product of the real numbers  $r$  and  $a$  and  $r$  and  $b$ , respectively. Then  $\cdot$  combines any real number with any member of  $\mathbb{R}^2$  to produce a unique member of  $\mathbb{R}^2$ . This is the so-called standard scalar product on the vector space  $\mathbb{R}^2$ .

## Example

Let  $S$  be the set of all  $2 \times 2$  matrices with real coefficients,

# Scalar Products

## Example

Let  $S = \mathbb{R}^2 = \{(a, b) : a, b \in \mathbb{R}\}$  (that is,  $S$  is the set “ $\mathbb{R}^2$ ” of ordered pairs of real numbers; you should have worked with this set in calculus and a first course in linear algebra). Now define  $\cdot$  by  $r \cdot (a, b) = (ra, rb)$ , where  $ra$  and  $rb$  are just the usual product of the real numbers  $r$  and  $a$  and  $r$  and  $b$ , respectively. Then  $\cdot$  combines any real number with any member of  $\mathbb{R}^2$  to produce a unique member of  $\mathbb{R}^2$ . This is the so-called standard scalar product on the vector space  $\mathbb{R}^2$ .

## Example

Let  $S$  be the set of all  $2 \times 2$  matrices with real coefficients, and define

$$r \cdot \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} ra & rb \\ rc & rd \end{bmatrix}.$$

# Scalar Products

## Example

Let  $S = \mathbb{R}^2 = \{(a, b) : a, b \in \mathbb{R}\}$  (that is,  $S$  is the set “ $\mathbb{R}^2$ ” of ordered pairs of real numbers; you should have worked with this set in calculus and a first course in linear algebra). Now define  $\cdot$  by  $r \cdot (a, b) = (ra, rb)$ , where  $ra$  and  $rb$  are just the usual product of the real numbers  $r$  and  $a$  and  $r$  and  $b$ , respectively. Then  $\cdot$  combines any real number with any member of  $\mathbb{R}^2$  to produce a unique member of  $\mathbb{R}^2$ . This is the so-called standard scalar product on the vector space  $\mathbb{R}^2$ .

## Example

Let  $S$  be the set of all  $2 \times 2$  matrices with real coefficients, and define

$$r \cdot \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} ra & rb \\ rc & rd \end{bmatrix}.$$

Then this is also a perfectly good scalar product on the set  $S = M_{2 \times 2}(\mathbb{R})$  of  $2 \times 2$  real matrices.

# Scalar Products

## Example

Let  $S = \mathbb{R}^2 = \{(a, b) : a, b \in \mathbb{R}\}$  (that is,  $S$  is the set “ $\mathbb{R}^2$ ” of ordered pairs of real numbers; you should have worked with this set in calculus and a first course in linear algebra). Now define  $\cdot$  by  $r \cdot (a, b) = (ra, rb)$ , where  $ra$  and  $rb$  are just the usual product of the real numbers  $r$  and  $a$  and  $r$  and  $b$ , respectively. Then  $\cdot$  combines any real number with any member of  $\mathbb{R}^2$  to produce a unique member of  $\mathbb{R}^2$ . This is the so-called standard scalar product on the vector space  $\mathbb{R}^2$ .

## Example

Let  $S$  be the set of all  $2 \times 2$  matrices with real coefficients, and define

$$r \cdot \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} ra & rb \\ rc & rd \end{bmatrix}.$$

Then this is also a perfectly good scalar product on the set  $S = M_{2 \times 2}(\mathbb{R})$  of  $2 \times 2$  real matrices.

# Scalar Products

## Example

Let  $S = \mathbb{R}$  and define  $r \cdot x = r + x$ .

# Scalar Products

## Example

Let  $S = \mathbb{R}$  and define  $r \cdot x = r + x$ . Then  $\cdot$  is the usual addition on  $\mathbb{R}$ :

# Scalar Products

## Example

Let  $S = \mathbb{R}$  and define  $r \cdot x = r + x$ . Then  $\cdot$  is the usual addition on  $\mathbb{R}$ : if we “combine” any real number  $r$  with any real number  $x$  (remember that  $S$  and  $\mathbb{R}$  are the same in this example),

# Scalar Products

## Example

Let  $S = \mathbb{R}$  and define  $r \cdot x = r + x$ . Then  $\cdot$  is the usual addition on  $\mathbb{R}$ : if we “combine” any real number  $r$  with any real number  $x$  (remember that  $S$  and  $\mathbb{R}$  are the same in this example), we get a uniquely determined member of  $S$ .

# Scalar Products

## Example

Let  $S = \mathbb{R}$  and define  $r \cdot x = r + x$ . Then  $\cdot$  is the usual addition on  $\mathbb{R}$ : if we “combine” any real number  $r$  with any real number  $x$  (remember that  $S$  and  $\mathbb{R}$  are the same in this example), we get a uniquely determined member of  $S$ .

## Example

Let  $S$  be the set  $\mathbb{N}$  of non-negative integers.

# Scalar Products

## Example

Let  $S = \mathbb{R}$  and define  $r \cdot x = r + x$ . Then  $\cdot$  is the usual addition on  $\mathbb{R}$ : if we “combine” any real number  $r$  with any real number  $x$  (remember that  $S$  and  $\mathbb{R}$  are the same in this example), we get a uniquely determined member of  $S$ .

## Example

Let  $S$  be the set  $\mathbb{N}$  of non-negative integers. Define  $\cdot$  by  $r \cdot n = -n$ .

# Scalar Products

## Example

Let  $S = \mathbb{R}$  and define  $r \cdot x = r + x$ . Then  $\cdot$  is the usual addition on  $\mathbb{R}$ : if we “combine” any real number  $r$  with any real number  $x$  (remember that  $S$  and  $\mathbb{R}$  are the same in this example), we get a uniquely determined member of  $S$ .

## Example

Let  $S$  be the set  $\mathbb{N}$  of non-negative integers. Define  $\cdot$  by  $r \cdot n = -n$ . Is  $\cdot$  a scalar product on  $S$ ?

# Scalar Products

## Example

Let  $S = \mathbb{R}$  and define  $r \cdot x = r + x$ . Then  $\cdot$  is the usual addition on  $\mathbb{R}$ : if we “combine” any real number  $r$  with any real number  $x$  (remember that  $S$  and  $\mathbb{R}$  are the same in this example), we get a uniquely determined member of  $S$ .

## Example

Let  $S$  be the set  $\mathbb{N}$  of non-negative integers. Define  $\cdot$  by  $r \cdot n = -n$ . Is  $\cdot$  a scalar product on  $S$ ?

*Solution.* No,  $\cdot$  is not a scalar product on  $S$ .

# Scalar Products

## Example

Let  $S = \mathbb{R}$  and define  $r \cdot x = r + x$ . Then  $\cdot$  is the usual addition on  $\mathbb{R}$ : if we “combine” any real number  $r$  with any real number  $x$  (remember that  $S$  and  $\mathbb{R}$  are the same in this example), we get a uniquely determined member of  $S$ .

## Example

Let  $S$  be the set  $\mathbb{N}$  of non-negative integers. Define  $\cdot$  by  $r \cdot n = -n$ . Is  $\cdot$  a scalar product on  $S$ ?

*Solution.* No,  $\cdot$  is not a scalar product on  $S$ . The issue is not uniqueness: for any real number  $r$  and non-negative integer  $n$ ,  $r \cdot n = -n$

# Scalar Products

## Example

Let  $S = \mathbb{R}$  and define  $r \cdot x = r + x$ . Then  $\cdot$  is the usual addition on  $\mathbb{R}$ : if we “combine” any real number  $r$  with any real number  $x$  (remember that  $S$  and  $\mathbb{R}$  are the same in this example), we get a uniquely determined member of  $S$ .

## Example

Let  $S$  be the set  $\mathbb{N}$  of non-negative integers. Define  $\cdot$  by  $r \cdot n = -n$ . Is  $\cdot$  a scalar product on  $S$ ?

*Solution.* No,  $\cdot$  is not a scalar product on  $S$ . The issue is not uniqueness: for any real number  $r$  and non-negative integer  $n$ ,  $r \cdot n = -n$  (and not, for example,  $-n$  AND  $-n + 1$  AND  $-n + 3$ ).

# Scalar Products

## Example

Let  $S = \mathbb{R}$  and define  $r \cdot x = r + x$ . Then  $\cdot$  is the usual addition on  $\mathbb{R}$ : if we “combine” any real number  $r$  with any real number  $x$  (remember that  $S$  and  $\mathbb{R}$  are the same in this example), we get a uniquely determined member of  $S$ .

## Example

Let  $S$  be the set  $\mathbb{N}$  of non-negative integers. Define  $\cdot$  by  $r \cdot n = -n$ . Is  $\cdot$  a scalar product on  $S$ ?

*Solution.* No,  $\cdot$  is not a scalar product on  $S$ . The issue is not uniqueness: for any real number  $r$  and non-negative integer  $n$ ,  $r \cdot n = -n$  (and not, for example,  $-n$  AND  $-n + 1$  AND  $-n + 3$ ). The issue is that  $r \cdot n$  is almost never a member of  $\mathbb{N}$ .

# Scalar Products

## Example

Let  $S = \mathbb{R}$  and define  $r \cdot x = r + x$ . Then  $\cdot$  is the usual addition on  $\mathbb{R}$ : if we “combine” any real number  $r$  with any real number  $x$  (remember that  $S$  and  $\mathbb{R}$  are the same in this example), we get a uniquely determined member of  $S$ .

## Example

Let  $S$  be the set  $\mathbb{N}$  of non-negative integers. Define  $\cdot$  by  $r \cdot n = -n$ . Is  $\cdot$  a scalar product on  $S$ ?

*Solution.* No,  $\cdot$  is not a scalar product on  $S$ . The issue is not uniqueness: for any real number  $r$  and non-negative integer  $n$ ,  $r \cdot n = -n$  (and not, for example,  $-n$  AND  $-n + 1$  AND  $-n + 3$ ). The issue is that  $r \cdot n$  is almost never a member of  $\mathbb{N}$  (it only is if  $n = 0$ ).

# Scalar Products

## Example

Let  $S = \mathbb{R}$  and define  $r \cdot x = r + x$ . Then  $\cdot$  is the usual addition on  $\mathbb{R}$ : if we “combine” any real number  $r$  with any real number  $x$  (remember that  $S$  and  $\mathbb{R}$  are the same in this example), we get a uniquely determined member of  $S$ .

## Example

Let  $S$  be the set  $\mathbb{N}$  of non-negative integers. Define  $\cdot$  by  $r \cdot n = -n$ . Is  $\cdot$  a scalar product on  $S$ ?

*Solution.* No,  $\cdot$  is not a scalar product on  $S$ . The issue is not uniqueness: for any real number  $r$  and non-negative integer  $n$ ,  $r \cdot n = -n$  (and not, for example,  $-n$  AND  $-n + 1$  AND  $-n + 3$ ). The issue is that  $r \cdot n$  is almost never a member of  $\mathbb{N}$  (it only is if  $n = 0$ ). For example,  $\sqrt{2} \cdot 4 = -4$ , but  $-4 \notin \mathbb{N}$ . □